


Provider Quarantine & Decon Procedures for Suspected COVID-19 Cases

Providers will carry out the following actions when in contact with patient(s) who have positive screening for Covid-19:

- Contact County-10 on duty who will provide guidance based on VDH Risk Assessment Tool (see attached)



The following steps SHALL be followed for movement of affected unit from hospital/scene:

1. All providers will don a gown, rubber exam gloves, N95 mask and some sort of eye protection.
2. Drive unit to rear parking lot of the New London station.
3. DC Johnston, BC Moore, or designee will meet driver to take ownership of unit and start the decon process.
4. Providers will be instructed to remove PPE and dispose of it in bin provided.
5. If further provider decon is necessary, shower facility at New London station will be used.

Coronavirus Disease 2019 (COVID-19) VDH Healthcare Personnel Risk Assessment Tool

Epidemiologic Risk Classification for Asymptomatic Healthcare Personnel Following Exposure to Patients with COVID19 or their Secretions/Excretions in a Healthcare Setting, and their Associated Monitoring and Work Restriction Recommendations

Epidemiologic risk factors	Exposure category	Recommended Monitoring for COVID-19 (until 14 days after last potential exposure)	Work Restrictions for Asymptomatic HCP
Prolonged close contact with a COVID-19 patient who was wearing a facemask (i.e., source control)			
HCP PPE: None	Medium	Active	Exclude from work for 14 days after last exposure
HCP PPE: Not wearing a facemask or respirator	Medium	Active	Exclude from work for 14 days after last exposure
HCP PPE: Not wearing eye protection	Low	Self with delegated supervision	None
HCP PPE: Not wearing gown or gloves ^a	Low	Self with delegated supervision	None
HCP PPE: Wearing all recommended PPE (except wearing a facemask instead of a respirator)	Low	Self with delegated supervision	None
Prolonged close contact with a COVID-19 patient who was not wearing a facemask (i.e., no source control)			
HCP PPE: None		Active	Exclude from work for 14 days after last exposure
HCP PPE: Not wearing a facemask or respirator		Active	Exclude from work for 14 days after fast exposure
HCP PPE: Not wearing eye protection ^b	Medium	Active	Exclude from work for 14 days after last exposure
HCP PPE: Not wearing gown or gloves ^{1b}	Low	Self with delegated supervision	None
HCP PPE: Wearing all recommended PPE (except wearing a facemask instead of a respirator) ^b	Low	Self with delegated supervision	None

HCP=healthcare personnel; PPE=personal protective equipment ^aThe risk category for these rows would be elevated by one level if HCP had extensive body contact with the patients (e.g., rolling the patient). ^bThe risk category for these rows would be elevated by one level if HCP performed or were present for a procedure likely to generate higher concentrations of respiratory secretions or aerosols (e.g., cardiopulmonary resuscitation, intubation, extubation, bronchoscopy, nebulizer therapy, sputum induction). For example, HCP who were wearing a gown, gloves, eye protection and a facemask (instead of a re5pirator) during an aerosolgenerating procedure would be considered to have a medium-risk exposure.